


Solutions to exam 2019-01-09

1a) $f(t) = \pi^2 - t^2 \quad -\pi \leq t \leq \pi$



Even fcn $\Rightarrow b_n = 0$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} (\pi^2 - t^2) \cos(nt) dt = \frac{2}{\pi} \int_0^{\pi} (\pi^2 - t^2) \cos(nt) dt =$$

$$\stackrel{n \geq 0}{=} \frac{2}{\pi} \left[(\pi^2 - t^2) \frac{\sin(nt)}{n} \right]_0^{\pi} - \frac{2}{\pi} \int_0^{\pi} -2t \frac{\sin(nt)}{n} dt$$

$$= \frac{4}{\pi n} \int_0^{\pi} t \sin(nt) dt = -\frac{4}{\pi n} \left[t \frac{\cos nt}{n} \right]_0^{\pi} +$$

$$+ \frac{4}{\pi n} \int_0^{\pi} \frac{\cos nt}{n} dt = -\frac{4}{\pi n^2} \pi \cos n\pi = \frac{4}{n^2} (-1)^{n+1}$$

$$\left[\frac{\sin nt}{n^2} \right]_0^{\pi} = 0$$

$$a_0 = \frac{2}{\pi} \int_0^{\pi} (\pi^2 - t^2) dt = \frac{2}{\pi} \left[\pi^2 t - \frac{t^3}{3} \right]_0^{\pi} = \frac{2}{\pi} \left(\pi^3 - \frac{\pi^3}{3} \right) =$$
$$= \frac{4\pi^2}{3}$$

$$\text{So } f(t) \sim \frac{2\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4}{n^2} (-1)^{n+1} \cos(nt)$$

1b) Since f is continuous and the generalised one-sided derivatives exist the Fourier series converges to f everywhere

1c) Take $t=0$. Then

$$\pi^2 = \frac{2\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4}{n^2} (-1)^n$$

$$\Rightarrow \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{\pi^2}{12}$$

1d) Parseval: $\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)|^2 dt = \frac{|a_0|^2}{4} + \frac{1}{2} \sum_{n=1}^{\infty} |a_n|^2 + |b_n|^2$

$\uparrow$
 $=0$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} (\pi^2 - t^2)^2 dt = \frac{1}{\pi} \int_0^{\pi} \pi^4 - 2t^2\pi^2 + t^4 dt =$$

$$= \frac{1}{\pi} \left[\pi^4 t - \frac{2}{3} t^3 \pi^2 + \frac{1}{5} t^5 \right]_0^{\pi} = \frac{1}{\pi} \left(\pi^5 - \frac{2}{3} \pi^5 + \frac{1}{5} \pi^5 \right)$$

$$= \pi^4 \left(\frac{15}{15} - \frac{10}{15} + \frac{3}{15} \right) = \frac{8}{15} \pi^4$$

Hence $\frac{8}{15} \pi^4 = \frac{4\pi^4}{9} + \frac{1}{2} \sum_{n=1}^{\infty} \frac{16}{n^4}$

$$\Rightarrow 8 \sum_{n=1}^{\infty} \frac{1}{n^4} = \left(\frac{8}{15} - \frac{4}{9} \right) \pi^4$$

$$\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^2} = \left(\frac{1}{15} - \frac{1}{18} \right) \pi^4 = \left(\frac{6}{90} - \frac{5}{90} \right) \pi^4 = \frac{\pi^4}{90}$$

$$2) \quad -u''(x) + u(x) = f(x)$$

Take the Fourier transform and use F.7:

$$\xi^2 \hat{u}(\xi) + \hat{u}(\xi) = \hat{f}(\xi)$$

$$\Rightarrow \hat{u}(\xi) = \frac{\hat{f}(\xi)}{1 + \xi^2}$$

Since $\hat{f}(\xi)$ is bounded we have $\frac{\hat{f}(\xi)}{1 + \xi^2} \in L^1(\mathbb{R})$ and inversion formula yields

$$u(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{f}(\xi)}{1 + \xi^2} e^{ix\xi} d\xi$$

$$3) \quad \text{Rewrite } f(x) = e^{-x^2} = e^{-(\sqrt{2}x)^2/2}$$

$$\text{Then } \hat{f}(\xi) = \frac{1}{\sqrt{2}} \sqrt{2\pi} e^{-\left(\frac{\xi}{\sqrt{2}}\right)^2/2} = \sqrt{\pi} e^{-\xi^2/4}$$

$$\text{By F.9 } \hat{g}(\xi) = (\hat{f}(\xi))^2 = \pi e^{-\xi^2/2} = \sqrt{\frac{\pi}{2}} \sqrt{2\pi} e^{-\xi^2/2}$$

$$\text{So } g(x) = \sqrt{\frac{\pi}{2}} e^{-x^2/2}$$

4) We use G-S to construct an ON-basis from the vectors

$f_0(x) = 1, f_1(x) = x$ in the inner product

$$\langle f, g \rangle := \int_0^{\infty} f(x) g(x) e^{-x} dx.$$

$$g_0(x) = f_0(x) = 1, \quad \|g_0\|^2 = \int_0^{\infty} |g_0(x)|^2 e^{-x} dx = \int_0^{\infty} e^{-x} dx = \left[-e^{-x} \right]_0^{\infty} = 1$$

$$e_0(x) = \frac{g_0(x)}{\|g_0\|} = 1$$

$$g_1(x) = f_1(x) - \langle e_0, f_1 \rangle e_0(x)$$

We have $\langle e_0, f_1 \rangle = \int_0^{\infty} x e^{-x} dx = \left[-x e^{-x} \right]_0^{\infty} + \underbrace{\int_0^{\infty} e^{-x} dx}_{1 \text{ (from before)}} = 1,$

so $g_1(x) = x - 1$. $\|g_1\|^2 = \int_0^{\infty} (x-1)^2 e^{-x} dx =$
 $= \int_0^{\infty} x^2 e^{-x} dx - 2 \underbrace{\int_0^{\infty} x e^{-x} dx}_{1 \text{ (from before)}} + \underbrace{\int_0^{\infty} e^{-x} dx}_{1} = \int_0^{\infty} x^2 e^{-x} dx - 1$

Hence we need to calculate

$$\int_0^{\infty} x^2 e^{-x} dx = \left[-x^2 e^{-x} \right]_0^{\infty} + \underbrace{\int_0^{\infty} 2x e^{-x} dx}_{2 \text{ (from before)}} = 2$$

So $\|g_1\|^2 = 1$ and $e_1(x) = g_1(x) = x - 1$.

The coefficients d_0, d_1 minimising

$$\int_0^{\infty} |c_0 e_0(x) + c_1 e_1(x) - \sin x|^2 e^{-x} dx$$

are $d_0 = \langle \sin, e_0 \rangle$, $d_1 = \langle \sin, e_1 \rangle$. To calculate these, it is useful to find the primitive function of $e^{-x} \sin x$, so let's do that first.

$$\int e^{-x} \sin x dx = -e^{-x} \cos x - \int e^{-x} \cos x dx$$

$$= -e^{-x} \cos x - \left(e^{-x} \sin x + \int e^{-x} \sin x dx \right) :$$

↑
Move this to the other side

$$\Rightarrow 2 \int e^{-x} \sin x dx = -e^{-x} (\cos x + \sin x)$$

$$\Rightarrow \int e^{-x} \sin x dx = -\frac{1}{2} e^{-x} (\cos x + \sin x)$$

$$\text{Now } d_0 = \int_0^{\infty} \sin x e^{-x} dx = -\frac{1}{2} \left[e^{-x} (\cos x + \sin x) \right]_0^{\infty} =$$

$$= \frac{1}{2}$$

$$d_1 = \int_0^{\infty} (x-1) \sin x e^{-x} dx = \int_0^{\infty} x \sin x e^{-x} dx - \underbrace{\int_0^{\infty} \sin x e^{-x} dx}_{1/2}$$

We have using IBP

$$\int_0^{\infty} x \sin x e^{-x} dx = \left[-\frac{1}{2} x e^{-x} (\sin x + \cos x) \right]_0^{\infty} + \underbrace{\frac{1}{2} \int_0^{\infty} e^{-x} \sin x dx}_{\frac{1}{2}}$$

$$+ \frac{1}{2} \int_0^{\infty} e^{-x} \cos x dx$$

$$\int_0^{\infty} e^{-x} \cos x dx = \frac{1}{2} \int_0^{\infty} (e^{(i-1)x} + e^{-(i+1)x}) dx = \frac{1}{2} \left[\frac{e^{(i-1)x}}{i-1} - \frac{e^{-(i+1)x}}{i+1} \right]_0^{\infty} =$$

$$= -\frac{1}{2} \left(\frac{1}{i-1} - \frac{1}{i+1} \right) = -\frac{1}{2} \frac{i+1-(i-1)}{-1-1} = \frac{1}{4} \cdot 2 = \frac{1}{2}$$

$$\text{Hence } \int_0^{\infty} x \sin x e^{-x} dx = \frac{1}{2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{2}$$

and

$$d_1 = \frac{1}{2} - \frac{1}{2} = 0.$$

The polynomial minimising the integral

$$d_0 e_0(x) + d_1 e_1(x) = \frac{1}{2} \cdot 1 + 0 \cdot (x-1) = \frac{1}{2} \cdot 1 + 0 \cdot x, \text{ so}$$

$$c_1 = \frac{1}{2}, \quad c_2 = 0$$

5) We homogenise using the ansatz $u(x,t) = v(x,t) + Ax + B$

$$BC \Rightarrow -\pi = v(0,t) + B, \quad 2\pi = v(\pi,t) + A\pi + B$$

This suggests $B = -\pi$, $A\pi = 3\pi \Rightarrow A = 3$

We also obtain $v(x,0) = u(x,0) - Ax - B = \pi - 3x$

Hence

$$\begin{cases} v_t = v_{xx} \\ v(0,t) = v(\pi,t) = 0 \\ v(x,0) = \pi - 3x \end{cases}$$

which is a homogeneous problem.

Now make the ansatz $v(x,t) = X(x)T(t)$.

Then $XT' = X''T$. Divide by XT on both sides so

$$\frac{T'}{T} = \frac{X''}{X} = -\lambda$$

↑ because none of LHS or RHS can be dependent on x or t

We start with the X equation and consider 3 cases?

Case 1 $\lambda < 0$: $X(x) = Ae^{\sqrt{-\lambda}x} + Be^{-\sqrt{-\lambda}x}$

Here $BC \Rightarrow \begin{cases} 0 = A + B \\ 0 = Ae^{\sqrt{-\lambda}\pi} + Be^{-\sqrt{-\lambda}\pi} \end{cases}$

which has the trivial solution $A=B=0$.

Case 2 $\lambda = 0$: $X(x) = Ax + B$

$BC \Rightarrow \begin{cases} 0 = A \cdot 0 + B \\ 0 = A\pi + B \end{cases} \Rightarrow A=B=0$

$$= -\frac{2}{\pi} \left((\pi - 2\pi) \begin{pmatrix} \cos(\pi) \\ \sin(\pi) \end{pmatrix} - \begin{pmatrix} \cos(0) \\ \sin(0) \end{pmatrix} \right) = \frac{1}{\pi} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Case 3 $\lambda > 0$: $X(x) = A \sin(\sqrt{\lambda} x) + B \cos(\sqrt{\lambda} x)$

$$X(0) = 0 \Rightarrow 0 = A \sin 0 + B \cos 0 \Rightarrow B = 0$$

$$X(\pi) = 0 \Rightarrow 0 = A \sin(\sqrt{\lambda} \pi)$$

$$\Rightarrow \sin(\sqrt{\lambda} \pi) = 0 \Rightarrow \sqrt{\lambda_n} \pi = n\pi \quad n=1, 2, \dots$$

$$\Rightarrow \lambda_n = n^2$$

Hence $X_n(x) = A_n \sin nx$

Now we consider T equation:

$$T'_n(t) = -n^2 T_n(t)$$

$$\Rightarrow T_n(t) = C_n e^{-n^2 t}$$

The full solution can be written as

$$v(x, t) = \sum_{n=1}^{\infty} X_n(x) T_n(t) = \sum_{n=1}^{\infty} A_n \sin(nx) e^{-n^2 t}$$

Consider the IC: $u(x, 0) = \pi - 3x$, i.e.

$$\sum_{n=1}^{\infty} A_n \sin(nx) = \pi - 3x \quad x \in (0, \pi)$$

Take $f(x) = \begin{cases} \pi - 3x & x \in (0, \pi) \\ -\pi - 3x & x \in (-\pi, 0] \end{cases}$ (odd extension)

$\{A_n\}$ are the Fourier coefficients of f

so $A_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx = \frac{2}{\pi} \int_0^{\pi} (\pi - 3x) \sin(nx) dx =$

$$= -\frac{2}{\pi} \left[(\pi - 3x) \frac{\cos(nx)}{n} \right]_0^{\pi} + \frac{2}{\pi} \int_0^{\pi} -3 \frac{\cos nx}{n} dx =$$

$$\underbrace{\left[-3 \frac{\sin nx}{n^2} \right]_0^{\pi}}_{=0}$$

$$= -\frac{2}{\pi} \left((\pi - 3\pi) \frac{\cos(n\pi)}{n} - \frac{\pi}{n} \right) = \frac{4(-1)^n + 2}{n}$$

Hence $v(x, t) = \sum_{n=1}^{\infty} \frac{4(-1)^n + 2}{n} \sin(nx) e^{-n^2 t}$

and therefore $u(x, t) = 3x - \pi + \sum_{n=1}^{\infty} \frac{4(-1)^n + 2}{n} \sin(nx) e^{-n^2 t}$

6) Taking Laplace transform gives

$$s^2 Y(s) - s y(0) - y'(0) + s(s Y(s) - y(0)) - 24 Y(s) = \frac{28}{s+1}$$

$$Y(s)(s^2 + 5s - 24) - 6s - 30 = \frac{28}{s+1}$$

$$Y(s) = \frac{\frac{28}{s+1} + \frac{6(s+5)(s+1)}{s+1}}{s^2 + 5s - 24}$$

$$\left(\text{obs: } s^2 + 5s - 24 = 0 \Rightarrow s = -\frac{5}{2} \pm \sqrt{\frac{25}{4} + \frac{96}{4}} = -\frac{5}{2} \pm \sqrt{\frac{121}{4}} \right. \\ \left. = -\frac{5}{2} \pm \frac{11}{2} \Rightarrow s_1 = -8, s_2 = 3 \Rightarrow s^2 + 5s - 24 = (s-3)(s+8) \right)$$

$$Y(s) = \frac{28 + 6(s+5)(s+1)}{(s+1)(s+8)(s-3)} = \frac{28 + 6s^2 + 36s + 30}{(s+1)(s+8)(s-3)} =$$

$$= \frac{6s^2 + 36s + 58}{(s+1)(s+8)(s-3)} \underset{\substack{\uparrow \\ \text{ansatz}}}{=} \frac{A}{s+1} + \frac{B}{s+8} + \frac{C}{s-3}$$

$$6s^2 + 36s + 58 = A(s+8)(s-3) + B(s+1)(s-3) + C(s+1)(s+8)$$

$$\text{Take } s = -8 \Rightarrow \underbrace{6 \cdot 8^2}_{384} - \underbrace{36 \cdot 8}_{288} + 58 = B(-7)(-11)$$

$$B = 2$$

$$\text{Take } s = -1 \Rightarrow \underbrace{6 - 36 + 58}_{28} = A(7)(-4)$$

$$\Rightarrow A = -1$$

$$\text{Take } s = 3 \Rightarrow \underbrace{6 \cdot 3^2}_{54} + \underbrace{36 \cdot 3}_{108} + 58 = C(4)(11)$$
$$C = 5$$

Hence $Y(s) = \frac{5}{s-3} + \frac{2}{s+8} - \frac{1}{s+1}$

Taking inverse Laplace transform gives
 $y(x) = 5e^{3x} + 2e^{-8x} - e^{-x}$

7) T is the convolution operator given by

$$Tf(x) = g * f(x) \quad \text{for } g(x) = \frac{2\sin x}{x}$$

Hence $\widehat{Tf}(\xi) = \widehat{g}(\xi) \widehat{f}(\xi)$

Using F.8 and F.10

$$\widehat{g}(\xi) = 2\pi \mathcal{K}_{[-1,1]}(-\xi) \stackrel{\text{even}}{=} 2\pi \mathcal{K}_{[-1,1]}(\xi)$$

Now Plancherel yields using $|\mathcal{K}_{[-1,1]}(\xi)| \leq 1$

$$\|Tf\|_{L^2}^2 = \frac{1}{2\pi} \|\widehat{Tf}\|_{L^2}^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} |2\pi \mathcal{K}_{[-1,1]}(\xi) \widehat{f}(\xi)|^2 d\xi$$

$$\leq 4\pi^2 \frac{1}{2\pi} \int_{-\infty}^{\infty} |\widehat{f}(\xi)|^2 d\xi = 4\pi^2 \int_{-\infty}^{\infty} |f(x)|^2 dx =$$

$$= 4\pi^2 \|f\|_{L^2}^2$$

Thus $\|Tf\|_{L^2} \leq 2\pi \|f\|_{L^2}$

8a) By definition $\hat{f}(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-int} dt$

IBP 2 times (allowed since $f \in C^2(\mathbb{T})$) yields

$$\begin{aligned} \hat{f}(n) &= \frac{1}{2\pi} \left[\underbrace{f(t) \frac{e^{-int}}{-in}}_{\in C(\mathbb{T}) \text{ and hence } 0} \right]_{-\pi}^{\pi} + \frac{1}{2\pi} \int_{-\pi}^{\pi} f'(t) \frac{e^{-int}}{in} dt \\ &= \frac{1}{2\pi} \left[\underbrace{f'(t) \frac{e^{-int}}{-(in)^2}}_{\in C(\mathbb{T}) \Rightarrow 0} \right]_{-\pi}^{\pi} + \frac{1}{2\pi} \int_{-\pi}^{\pi} f''(t) \frac{e^{-int}}{(in)^2} dt \end{aligned}$$

Now triangle inequality gives

$$|\hat{f}(n)| \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |f''(t)| \frac{1}{n^2} dt \leq \frac{C}{n^2} \quad \text{where}$$

$$C = \frac{1}{2\pi} \int_{-\pi}^{\pi} |f''(t)| dt$$

b) Yes! To prove this, use Weierstrass' M-test. The Fourier series is given by

$$\sum_{n=-\infty}^{\infty} \hat{f}(n) e^{int}$$

But since $|\hat{f}(n) e^{int}| \leq \frac{C}{n^2}$ and $\sum_{n \neq 0} \frac{C}{n^2} < \infty$

the M-test applies and the Fourier series is uniformly convergent.